



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

settlement of the western portion of Kansas will have a similar effect upon its rainfall; but it is not reasonable to expect that western Kansas will ever boast of a rainfall equal to that of eastern Kansas. So long as the eastern half of the state remains to the east of the meridian forming the western boundary of the Gulf of Mexico, the south winds will cause it to receive much larger supplies of vapor, for condensation into rain, than will be received by the western half of the state, which lies beyond the immediate track of the vapor-laden winds. It must be remembered that climatic changes are exceedingly gradual; and a rain deficiency or excess for a single year, or for two or three years in succession, must not be considered as invalidating the law of general averages. Neither should the fact that the rainfall, upon the whole, is increasing, induce settlers to break land in the western third of Kansas with the expectation of successfully raising the same crops as in eastern Kansas. Such settlers will surely be disappointed. It is even doubtful if paying crops of any kind can ever be continuously produced in that region. With an average before settlement of about fifteen inches per annum, the same percentage of increase as has been made in eastern Kansas in thirty years would give an annual amount of less than eighteen inches, — a quantity entirely inadequate to maintain successful agriculture.

AMERICAN SOCIETY FOR PSYCHICAL RESEARCH.

At a meeting held in Boston, Sept 23, to consider the advisability of the formation of a society for psychical research in America, the whole matter was placed in the hands of a committee of nine, consisting of Dr. G. Stanley Hall of Johns Hopkins university; Prof. E. C. Pickering, director of the Harvard college observatory; Dr. H. P. Bowditch and Dr. C. S. Minot, of the Harvard medical school; Mr. S. H. Scudder, president, and Professor Alpheus Hyatt, curator, of the Boston society of natural history; Professor William James of Harvard college; Professor William Watson of Boston; and Mr. N. D. C. Hodges of Cambridge. This committee held a number of meetings during the months of October and November, and issued an invitation to a number of scientific men throughout the country to join in a society under a constitution upon which they had decided. To this invitation there were favorable replies from about eighty.

The first meeting of the society was held in Boston on the 18th of December. Under the constitution the conduct of the society is placed in the hands of a council of twenty-one, seven to be chosen each year, to hold office three years. Of this council, there were elected at this first meeting, fifteen: Prof. G. Stanley Hall, Prof. George S. Fullerton, Dr. William James, Prof. E. C. Pickering, for three years; Professor Simon Newcomb, Dr. C. S. Minot, Dr. H. P. Bowditch, Mr. N. D. C. Hodges, for two years; Prof. George F. Barker, Mr. S. H. Scudder, Rev. C. C. Everett, Mr.

Morefield Storey, Professor John Trowbridge, Professor William Watson, Professor Alpheus Hyatt, for one year.

The sub-committee on work made an informal report, and has since issued a circular to members, asking for volunteers on the investigating committees and for information regarding promising subjects for investigation, such as mediums, mind-readers, mesmeric subjects, etc.

The society adjourned to meet on the ninth day of January.

THE NATURAL BRIDGE OF VIRGINIA.¹

DURING a recent trip to Virginia (Oct. 2 to 6) I visited the Natural Bridge; and although in possession of the guide-book of the locality (edition of 1884), and the admirable articles published by Major Jed. Hotchkiss in *The Virginias*, I failed to obtain certain information relating to the bridge, which would be of special interest to the topographer and geologist. Some of the observations which I made, although of a general character, may be of interest.

The bridge is undoubtedly the remnant of the top of a cave which was probably formed long before the Luray cavern, which is excavated out of the same lower Silurian limestone formation. The bridge seems to be located in the centre of a gentle basin or synclinal in the strata, which may account for the roof of the ancient cavern being left at this special point. The height of the bridge has evidently been much augmented by a lowering of the bed of Cedar Creek through the agency of chemical and mechanical erosion after the destruction of the original cavern. The height of the original cavity, at the point where the bridge now exists, was in consequence very much less than the present height of the intrados of the bridge-arch.

The elevation of the railroad-track at Natural Bridge station, on the Shenandoah valley railroad, is seven hundred and sixty feet above ocean-level; and the elevation of Cedar Creek, under the north face of the bridge-arch, is nine hundred and fifteen feet, as determined by two independent lines of barometric levels which I ran between the railroad-station and the bridge.

The height of the crown of the arch on the north side, at the 'Lookout Point,' is one hundred and eighty-eight feet above the creek, measured with a cotton twine, which was the only line of the required length which could be obtained. The same height measured by the barometer (Short & Mason aluminum aneroid) was determined as one hundred and eighty-six feet. Neither of these methods of measurement is sufficiently exact to permit of a final statement, but the results are of interest in the absence of more definite data.

The thickness of the arch under the crown on the north side is approximately forty-six feet, and on the south side thirty-six feet.

¹ Read before the American philosophical society, Oct. 17, 1884, by CHARLES A. ASHBURNER.